

HOPS

This sheet lists some pesticides used on hops. The health symptoms listed may be caused by things other than pesticides. If you feel sick after exposure to pesticides, contact a doctor or clinic.



Year 2002

Pesticide (Brand Name) highly toxic	Product Use Changes Read the Label DO NOT ENTER TREATED AREAS FOR: Reentry intervals will vary by product	Common Symptoms of Acute Poisoning
Abamectin (Agri-Mek)	1/2 Day / 12 hours	NOSE EAR, OR THROAT IRRITATION BREATHING PROBLEMS STOMACH ACHE WEAKNESS
Bifenthrin (Brigade)	1 Day	STOMACH DIZZINESS VOMITING IRRITATION
Endothall (Des-I-Cate)	2 Days	NOSE EAR OR THROAT IRRITATION SKIN IRRITATION VOMITING WEAKNESS VISION PROBLEMS
Imidacloprid (Provado)	1/2 Day / 12 hours	NOSE, EAR, OR THROAT IRRITATION IRRITATION
Metalaxyl / Mefenozam (Ridomil Gold)	2 Days	NOSE EAR OR THROAT IRRITATION SKIN IRRITATION
Paraquat (Gramoxone Extra)	1 Day	NOSE EAR, OR THROAT IRRITATION IRRITATION
Propargite (Omite)	7 Days	NOSE EAR OR THROAT IRRITATION IRRITATION PROBLEMS HEADACHES
Sulfur	1 Day	NOSE EAR OR THROAT IRRITATION IRRITATION IRRITATION IRRITATION

If you are exposed to a pesticide spray or drift:

Leave the area



Tell your employer



Immediately wash exposed skin with soap and lots of water




If you get pesticides in your eyes, rinse with clean water for 15 minutes

Change into clean clothes, wash work clothes separately from other clothes



WSDA: Toll Free **1-877-301-4555** L&I Toll Free **1-800-423-7233**



Leave the area



Tell your employer you feel sick



Go to a doctor or clinic

Have someone go with you

Tell doctor the pesticide name. Your employer must provide this information





Emergency Pesticide Information:
WA Poison Control Center 1-800-222-1222
Pesticide Related Health Concerns:
DOH (Olympia) 360-236-3360